

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/225072572>

Overnutrition and associated factors among adults aged 20 years and above in fishing communities in the urban Cape Coast Metropolis, Ghana

Article in *Public Health Nutrition* · May 2012

DOI: 10.1017/S1368980012002698 · Source: PubMed

CITATIONS

16

READS

901

5 authors, including:



Kingsley Pereko

University of Cape Coast

18 PUBLICATIONS 32 CITATIONS

[SEE PROFILE](#)



Jacob Setorglo

University of Cape Coast

4 PUBLICATIONS 21 CITATIONS

[SEE PROFILE](#)



Emmanuel Achampong

University of Cape Coast

25 PUBLICATIONS 69 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Non-Alcoholic Fatty Liver Disease, Molecular Bases, Prevention and Treatment [View project](#)



Cloud computing and Electronic Health Records in Developing Countries [View project](#)

Overnutrition and associated factors among adults aged 20 years and above in fishing communities in the urban Cape Coast Metropolis, Ghana

Kingsley KA Pereko^{1,*}, Jacob Setorglo², William B Owusu³, Joyce M Tiweh⁴ and Emmanuel K Achampong⁵

¹Department of Community Medicine, School of Medical Sciences, University of Cape Coast, Cape Coast, Ghana: ²Department of Biochemistry, School of Medical Sciences, University of Cape Coast, Cape Coast, Ghana: ³Department of Nutrition and Food Science, University of Ghana, Accra, Ghana: ⁴Department of Ophthalmology Komfo Anokye Teaching Hospital, Kumasi, Ghana: ⁵Department of Medical Education, University of Cape Coast, Cape Coast, Ghana

Submitted 16 December 2010: Final revision received 16 February 2012: Accepted 17 April 2012: First published online 29 May 2012

Abstract

Objective: The study aimed to highlight the determinants of overnutrition (overweight plus obesity) in fishing communities and establish if these were the same as reported elsewhere in Ghana.

Design: Cross-sectional study.

Setting: The study was conducted in Idun, Ola and Duakor fishing communities in Cape Coast, Ghana.

Subjects: Adults (*n* 252) aged 20 to 50 years.

Results: Results showed that 32% of participants were overweight/obese (BMI ≥ 25.0 kg/m²). Participants' mean age was 31.7 (SD 1.0) years, they had 13.7 (SD 8.1) mean years of formal education, their median monthly income was \$US 7.4 (interquartile range \$US 3.3, 20.0) and their median daily energy intake was 7.3 (interquartile range 5.3, 9.8) MJ. Significant associations ($P < 0.05$) were found between BMI and gender, age, years of education, fat intake and marital status. Females were almost eight times more likely to be overweight/obese than males (adjusted OR = 7.7; 95% CI 3.6, 16.4). Persons aged ≥ 40 years were about six times more likely to be overweight/obese than those aged 20–29 years (adjusted OR = 6.1; 95% CI 2.6, 14.1). Married people were nearly three times more likely to be overweight/obese than singles (adjusted OR = 2.8; 95% CI 1.4, 5.7). People with more than 13 years of formal education (adjusted OR = 0.3; 95% CI 0.1, 0.9) and people with $>30\%$ fat contribution to daily energy intake (adjusted OR = 0.3; 95% CI 0.1, 0.6) had reduced odds of being overweight/obese.

Conclusions: Overnutrition was prevalent in the fishing communities and associated with factors such as age, gender, marital status, educational status and fat intake.

Keywords
Overnutrition
Macronutrient
Formal education
Energy

Overnutrition, a major public health issue of the developed world, is a growing menace in the developing world already burdened with undernutrition and infectious diseases⁽¹⁾. Gupta and Kockar⁽²⁾ indicate obesity to be one of the most neglected public health problems according to WHO. Hajian-Tilaki and Heidari⁽³⁾ state that obesity is an undesirable outcome of changing lifestyles and behaviours. Overnutrition is an emerging problem in segments of Sub-Saharan African society particularly where lifestyles have become urbanized and Westernized⁽⁴⁾. The issue of overnutrition has become very important as a result of its debilitating effect on health. Obesity increases the risk for many serious and morbid conditions, such as diabetes

mellitus, hypertension, dyslipidaemia, coronary artery diseases and some cancers^(1,4). Seidell⁽⁵⁾ found obesity to be 1.5–2.0 times higher in women than men from countries with low gross national product. Studies in adolescents have found an association of BMI with gender, ethnicity and food habits. In other studies, obesity has been found to be higher among married women than unmarried⁽⁶⁾. Socio-economic status has been found to have an inverse relationship with obesity in the developed world; however, the opposite is the case in the developing world⁽⁷⁾. Work by Amoah⁽⁸⁾ in urban and rural Accra, Ghana showed crude prevalences of overweight and obesity of 23.4% and 14.1%, respectively, among adults aged 25 years and above. Females recorded

higher rates than males. Obesity increased with age and the rate was found to be higher among urban high-class residence compared with low-class residence and in urban compared with rural locations. Individuals with higher educational status recorded higher prevalence than those with lower education. A study in Accra involving Ghanaian civil servants found obesity prevalence of 10% among males and 36% among females⁽⁹⁾. However, the subject of obesity in Ghana receives much less attention than it deserves despite the growing menace⁽¹⁰⁾. The present study aimed to highlight the determinants of overweight in three fishing communities to establish if these were the same as those reported elsewhere in Ghana. The specific objectives were to: (i) examine the participants' sociodemographic characteristics (such as gender, occupation, marital status and years of education); (ii) assess their BMI and the dietary intake; and (iii) determine the associations between sociodemographic characteristics, dietary intake and BMI.

Methodology

The present study was a cross-sectional one involving 252 adults aged 20 to 50 years from the fishing communities of Ola, Duakor and Idun of the Cape Coast metropolis in Ghana. These communities had a total population of 10 708, comprising 7375, 1039 and 2294, respectively⁽¹¹⁾. Given an adult population size of 10 708 in the three communities and an expected frequency of 23.4% urban adult overweight and obesity prevalence, with worst acceptable frequency of 28.6% at 95% confidence interval, a sample size of 249 was needed for the study (Epi Info version 3.5.1; Centers for Disease Control and Prevention, Atlanta, GA, USA). From the total population of the chosen communities, weighting was used to estimate the number of participants required from each community. Based on the weighting, eighty-one, eighty-four and eighty-seven participants were randomly chosen from Ola, Duakor and Idun, respectively. Healthy-looking adults who self-reported absence of any ailment were included in the study while pregnant women and ill persons were excluded. A multistage sampling procedure was used. This involved the random selection of three out of four fishing communities in the urban metropolis. Systematic random sampling was used to select the houses in the communities. If there was more than one household in the same house, simple random sampling was used to select a respondent. Data were collected on socio-demographic factors using a semi-structured questionnaire and on dietary intake using a 24 h recall questionnaire. The 24 h dietary recall was conducted on one weekend and two weekdays for each of the respondents.

Data on anthropometry were collected following a standard protocol by taking two repeated measures⁽¹²⁾. Weight and height measures were taken using a

UNISCALE (UNICEF electronic scale 890) and a UNICEF adult microtoise (UNICEF, Copenhagen, Denmark), respectively, which were calibrated to ensure accuracy of the measures. Weight was measured with the participants in light clothing, with the scale on a level stable surface, to the nearest 0.1 kg. Participants were weighed while standing unsupported, bare footed and looking forward, with their hands by their side, and were not holding any metallic or other object that could influence the SECA scale. Weight measurements were read only when the digital reader was stable. Height of the participants in bare feet was measured to the nearest 0.5 cm using a standard microtoise. In measuring height, the head bar of the microtoise was placed securely on the floor and the tape fully drawn. The free end was secured with a nail to a suitable vertical surface above the head bar. The head bar was then raised above the height of the participant, who was instructed to stand upright directly below the point of attachment and positioned such that the Frankfort plane of his/her head was horizontal. The participant was instructed to inspire and to hold his/her breath while the head bar was lowered to the crown of the head, and a direct height reading in centimetres was taken. This was all administered by the interviewer.

Using the WHO reference for BMI, participants were classified into the following categories: BMI < 18.5 kg/m² (thin), BMI = 18.5–24.9 kg/m² (normal weight), BMI = 25.0–29.9 kg/m² (overweight) and BMI ≥ 30.0 kg/m² (obese)⁽¹³⁾. In the present paper, 'overnutrition' is taken to include overweight and obesity (BMI ≥ 25.0 kg/m²). Occupation of the participants was classified into sedentary and manual. For the study purposes, those who by their occupation spend more than five out of eight working hours in the day moving or doing manual work were considered physically active, while sedentary participants were those who spend more than five out of the eight working hours in a day sitting or not doing manual work.

The data were entered and analysed using the IBM SPSS version 20.0 statistical software package (IBM Corp, Armonk, NY, USA). A nutrition database (ESHA Food Processor[®] 6.02⁽¹⁴⁾) and the Ghanaian food composition table⁽¹⁵⁾ were used in estimating nutrient intakes. Means, standard deviations and ranges were calculated for continuous variables, while frequencies and proportions were calculated for categorical variables. The associations between demographic factors, dietary intake and BMI were assessed using the χ^2 test and logistic regression technique. Differences in means among males and females were determined using the independent-samples *t* test.

Results

In all 252 adults participated in the present study, of whom 40.1% were male and 59.9% were female (Table 1). Participants' mean age was 31.7 (SD 1.0) years and they had

Table 1 Descriptive statistics (demographic characteristics, anthropometric measures) of the study population: adults aged 20 to 50 years from three fishing communities in the urban Cape Coast metropolis, Ghana

	Males (n 101)			Females (n 151)			Both sexes (n 252)		
	Mean	SD	Range	Mean	SD	Range	Mean	SD	Range
Age (years)	31.6	8.1	20.0–51.0	31.7	8.9	20–50	31.7	1.0	20–51
Years of education	8.4	4.4	0.0–22.0	7.8	4.4	0.0–19.0	13.7	8.1	0.0–22.0
Weight (kg)	65.9	8.2	48.2–89.0	65.4	14.8	39.0–111.8	65.6	12.6	39.0–111.8
Height (m)	1.7	0.1	1.0–2.0	1.6*	0.1	1.2–1.8	1.6	0.1	1.0–2.0
BMI (kg/m ²)	23.3	6.6	17.8–29.9	25.8*	6.1	18.7–27.1	24.8	6.4	17.8–29.9

*Mean values were significantly different from those of males (independent-samples *t* test): $P < 0.05$.

Table 2 Descriptive statistics (income, dietary intake) of the study population: adults aged 20 to 50 years from three fishing communities in the urban Cape Coast metropolis, Ghana

	Males (n 101)		Females (n 151)		Both sexes (n 252)	
	Median	IQR	Median	IQR	Median	IQR
Monthly income (\$US)	10.0	6.7, 22.2	6.7	3.3, 20.0	7.4	3.3, 20.0
Energy (kJ)	9.0	6.8, 11.2	6.5	4.7, 8.7	7.3	5.3, 9.8
% of energy from fat	22.0	14.0, 34.5	24.0	15.0, 36.6	23.0	14.3, 36.0
% of energy from protein	11.0	10.0, 14.0	12.0	10.0, 14.6	12.0	10.0, 14.0
% of energy from carbohydrate	67.0	53.5, 74.0	61.0	51.0, 71.0	62.0	52.0, 72.0

IQR, interquartile range.

13.7 (SD 8.1) mean years of formal education. Mean BMI of 24.8 (SD 6.4) kg/m² was recorded for both sexes. Females had a significantly higher mean BMI (25.8 kg/m²) than males (23.3 kg/m²).

The median monthly income of the participants was \$US 7.4 (interquartile range (IQR) \$US 3.3, 20.0; Table 2). The participants' median energy intake was 7.3 (IQR 5.3, 9.8) MJ and the median percentage contributions of macronutrients to daily energy intake were 23.0 (IQR 14.3, 36.0) % for fat, 12.0 (IQR 10.0, 14.0) % for protein and 62.0 (IQR 52.0, 72.0) % for carbohydrate (Table 2).

The results indicated an overweight/obesity prevalence of 32% (Table 3). More females (45.7%) than males (13.9%, $P < 0.05$) were found to be in the overweight/obese category. A significant difference in the proportion of overweight/obesity was found between married persons (20.3%) and singles (45.0%, $P < 0.05$; Table 3).

Females were almost eight times more likely to be overweight/obese than males (adjusted OR = 7.7; 95% CI 3.6, 16.4; Table 4). People aged ≥ 40 years were about six times more likely to be overweight/obese than those aged 20–29 years (adjusted OR = 6.1; 95% CI 2.6, 14.1; Table 4). Married people were almost three times more likely to be overweight/obese than singles (adjusted OR = 2.8; 95% CI 1.4, 5.7; Table 4). Compared with 13 years of education or less, participants with more than 13 years of education were 70.0% less likely to be overweight/obese (adjusted OR = 0.3; 95% CI 0.1, 0.9; Table 4). Participants with >30 % contribution to daily energy intake from fat were 70.0% less likely to be overweight/obese (adjusted OR = 0.3; 95% CI 0.1, 0.6; Table 4) than those with ≤ 30 % of daily energy intake from fat.

Discussion

In Ghana, studies show that overnutrition is gradually becoming an important issue following the increasing prevalence of overweight and obesity in most parts of the country. Information on the situation will be an important element for early management and prevention of overweight and obesity to avert its health-debilitating effects. Work by Amoah⁽⁸⁾ found an overweight prevalence of slightly over 23% in the urban population. The present study, however, found about 32% overnutrition prevalence among the study population in the urban fishing community. Factors that were found to be associated with this phenomenon were age, marital status, gender, years of formal education and dietary fat intake.

Age was found to be associated with overnutrition; the present study indicated 2.3 times increased odds of overweight/obesity among people aged 30–39 years compared with those aged 20–29 years and a 6.1 times increased likelihood for persons aged ≥ 40 years. This confirms van der Sande *et al.*'s⁽⁴⁾ study that recorded higher obesity prevalence among women aged >35 years. Biritwum *et al.*⁽¹⁰⁾ and Beltaifa *et al.*⁽¹⁶⁾ also found increasing obesity prevalence with increasing age. The present study found an association between marital status and overweight, with married persons being 2.8 times more likely to be overweight/obese compared with unmarried persons. This result affirms studies by Biritwum *et al.*⁽¹⁰⁾ and Rguibi and Belahsen⁽⁶⁾. Females were 7.7 times more likely to be overweight/obese than males in the present study and this confirms findings by van der Sande *et al.*⁽⁴⁾ and Biritwum *et al.*⁽¹⁰⁾ who reported similar

Table 3 Demographic characteristics and dietary intake according to overweight/obesity status: adults aged 20 to 50 years from three fishing communities in the urban Cape Coast metropolis, Ghana

	Not overweight/obese (BMI < 25.0 kg/m ²)		Overweight/obese (BMI ≥ 25.0 kg/m ²)	
	<i>n</i>	%	<i>n</i>	%
Total sample	169	67.1	83	32.9
Gender				
Male	87	86.1	14	13.9*
Female	82	54.3	69	45.7*
Monthly income (\$US)				
≤10	62	59.6	42	40.4
>10	60	65.2	32	34.2
Occupation				
Sedentary	79	63.2	46	36.8
Manual	90	70.9	37	29.1
Marital status				
Single	98	79.7	25	20.3*
Married	71	55.0	58	45.0*
Years of education				
≤13	139	64.1	78	35.9*
>13	29	85.3	5	14.7*
% contribution of fat to daily energy intake				
≤30	95	60.5	62	39.5*
>30	74	77.9	21	22.1*
% contribution of carbohydrate to daily energy intake				
≤60	38	67.9	18	32.1
>60	131	66.8	65	33.2
% contribution of protein to daily energy intake				
≤20	164	68.0	77	32.0
>20	5	45.5	6	54.5

*Proportions were significantly different from those in the not overweight/obese group (χ^2 test): $P < 0.05$.

Table 4 Logistic regression coefficients and 95% confidence intervals for predictor variables of overnutrition (BMI ≥ 25.0 kg/m²): adults aged 20 to 50 years from three fishing communities in the urban Cape Coast metropolis, Ghana

Variable	<i>R</i> ² (0.29)	
	OR	95% CI
Age (years)		
20–29	1.0 (ref.)	
30–39	2.3	1.1, 5.1
≥40	6.1	2.6, 14.1
Sex		
Male	1.0 (ref.)	
Female	7.7	3.6, 16.4
Years of formal education		
≤13	1.0 (ref.)	
>13	0.3	0.1, 0.9
Marital status		
Single	1.0 (ref.)	
Married	2.8	1.4, 5.7
% contribution of fat to daily energy intake		
≤30	1.0 (ref.)	
>30	0.3	0.1, 0.6

ref., reference category.

The regression coefficient for the entire model; $R^2 = 0.21$.

results. Females have a greater physiological propensity to become obese than males. Coupled with the sedentary occupations of women in the area, they have a greater chance to become overweight/obese than males who engage in energy-demanding activities such as fishing.

According to van der Sande *et al.*⁽⁴⁾, nothing can be done about the age and sex susceptibility but lifestyle modifications are possible. Contrary to Amoah's⁽⁸⁾ and Biritwum *et al.*'s⁽¹⁰⁾ finding of increasing obesity prevalence among persons with higher education, the present study found reduced odds of overweight/obesity among persons with higher education compared with those having basic or no formal education. This could be attributed to the fact that more highly educated persons are aware of the negative consequences of being overweight such as the development of chronic diseases, and hence take measures to prevent it. In their study of obesity among adults aged 20 to 70 years, Hajian-Tilaki and Heidari⁽³⁾ also found an inverse relationship between the risk of obesity and high level of education. Beltaifa *et al.*⁽¹⁶⁾ indicated a higher prevalence among women with intermediary educational status. Persons whose fat intake contributed >30% of daily energy intake were found less likely to be overweight/obese in the present study. This could possibly be attributed to the occupation they undertake. Males found to consume more fat probably undertake energy-demanding activities, hence accounting for the observation.

Conclusions

Overweight and obesity was prevalent in fishing communities in the Cape Coast urban metropolis, Ghana. Factors

such as age, gender, marital status, education and fat intake were associated with overnutrition in this community. Further prospective research is recommended for assessing the factors accounting for overweight and obesity. Health education on healthy lifestyles is recommended.

Acknowledgements

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors. There are no conflicts of interest. The authors' contributions are as follows: K.K.A.P., Principal Investigator; J.M.T., Co-Principal Investigator; J.S., project staff training, manuscript writing and review; W.B.O., supervisor, statistician; E.K.A., data entry, manuscript writing and review. The authors acknowledge the kind assistance and permission of the Cape Coast Metropolitan Health Directorate and Assembly Members of the three project communities. They also appreciate the participation of the Duakor, Idun and Ola fishing communities.

References

1. Martorell R, Khan LK, Hughes ML *et al.* (2000) Obesity in women from developing countries. *Eur J Clin Nutr* **54**, 247–252.
2. Gupta N & Kockar GK (2009) Dietary and socio-economic factors associated with obesity in Northern India population. *Internet J Health* **9**, 1.
3. Hajian-Tilaki KO & Heidari B (2007) Prevalence of obesity, central obesity and the associated factors in Urban population aged 20–70 years, in the North of Iran. A population-based study and regression approach. *Obes Rev* **8**, 3–10.
4. van der Sande MAB, Caesay SM, Milligan PJM *et al.* (2001) Obesity and undernutrition and cardiovascular risk factors in rural and urban Gambian communities. *Am J Public Health* **91**, 1641–1644.
5. Seidell JC (2005) Epidemiology of obesity. *Semin Vasc Med* **5**, 3–14.
6. Rguibi M & Belahsen R (2004) Obesity among urban Sahraoui women of South Morocco. *Ethin Dis* **14**, 542–547.
7. McLaren L (2007) Socioeconomic status and obesity. *Epidemiol Rev* **29**, 29–48.
8. Amoah AG (2003) Obesity in adult residents of Accra, Ghana. *Ethin Dis* **13**, 2 Suppl. 2, 29–101.
9. Addo J, Smeeth L & Leon DA (2009) Obesity in urban civil servants in Ghana. Association with pre-adult wealth and adult socio-economic status. *J Pubic Health* **123**, 365–370.
10. Biritwum RB, Gyapong J & Mensah G (2005) The epidemiology of obesity in Ghana. *Ghana Med J* **39**, 82–85.
11. Ghana Statistical Service (2002) *2000 Population and Housing Census. Special Report on 20 Largest Localities*. Accra: GSS.
12. Ismail M & Manandhar M (1999) *Better Nutrition for Elderly People. Assessment and Action*, p. 71. London: HelpAge International and London School of Hygiene and Tropical Medicine.
13. World Health Organization (2000) *Obesity Prevention and Managing Global Epidemic: Report of a WHO Consultation. WHO Technical Report Series* no. 894. Geneva: WHO.
14. Davison S & Mandible D (1994) *The Food Processor Plus 6.02*. Salem, OR: K.G. Dewey Lab, ESHA.
15. Tayie FAK & Lartey A (1999) *Nutrient Contents of Some Ghanaian Foods*. Accra: Nutrition and Food Science Department, University of Ghana.
16. Beltaifa L, Traissac P, El Ati J *et al.* (2009) Prevalence of obesity and associated socioeconomic factors among Tunisian women from different living environments. *Obes Rev* **10**, 145–153.